

Stochastic Algorithms for Solving Constrained Continuous Optimization Problems

Frank E. Curtis, Lehigh University

involving joint work with

Albert S. Berahas (U. of Michigan), **Xin Jiang** (Lehigh), **Vyacheslav Kungurtsev** (Czech TU),
Suyun Liu (Amazon), **Michael O'Neill** (UNC Chapel Hill), **Daniel P. Robinson** (Lehigh),
Qi Wang (Lehigh), **Baoyu Zhou** (Chicago Booth)


presented at

Department of Industrial and Manufacturing Engineering, Pennsylvania State University

November 1, 2023



Collaborators and references



- ▶ A. S. Berahas, F. E. Curtis, D. P. Robinson, and B. Zhou, “Sequential Quadratic Optimization for Nonlinear Equality Constrained Stochastic Optimization,” *SIAM Journal on Optimization*, 31(2):1352–1379, 2021.
- ▶ A. S. Berahas, F. E. Curtis, M. J. O’Neill, and D. P. Robinson, “A Stochastic Sequential Quadratic Optimization Algorithm for Nonlinear Equality Constrained Optimization with Rank-Deficient Jacobians,” to appear in *Mathematics of Operations Research*.
- ▶ F. E. Curtis, D. P. Robinson, and B. Zhou, “Inexact Sequential Quadratic Optimization for Minimizing a Stochastic Objective Subject to Deterministic Nonlinear Equality Constraints,” <https://arxiv.org/abs/2107.03512>.
- ▶ F. E. Curtis, M. J. O’Neill, and D. P. Robinson, “Worst-Case Complexity of an SQP Method for Nonlinear Equality Constrained Stochastic Optimization,” to appear in *Mathematical Programming*.
- ▶ F. E. Curtis, S. Liu, and D. P. Robinson, “Fair Machine Learning through Constrained Stochastic Optimization and an ϵ -Constraint Method,” to appear in *Optimization Letters*.
- ▶ F. E. Curtis, D. P. Robinson, and B. Zhou, “Sequential Quadratic Optimization for Stochastic Optimization with Deterministic Nonlinear Inequality and Equality Constraints,” <https://arxiv.org/abs/2302.14790>.
- ▶ F. E. Curtis, V. Kungurtsev, D. P. Robinson, and Q. Wang, “A Stochastic-Gradient-based Interior-Point Algorithm for Solving Smooth Bound-Constrained Optimization Problems,” <https://arxiv.org/abs/2304.14907>.

Outline

Motivation

Stochastic SQP

Extensions

Conclusion

Outline

Motivation

Stochastic SQP

Extensions

Conclusion

Constrained optimization (deterministic)

Consider

$$\begin{array}{ll} \min_{x \in \mathbb{R}^n} & f(x) \\ \text{s.t.} & c_{\mathcal{E}}(x) = 0 \\ & c_{\mathcal{I}}(x) \leq 0 \end{array}$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $c_{\mathcal{E}} : \mathbb{R}^n \rightarrow \mathbb{R}^{m_{\mathcal{E}}}$, and $c_{\mathcal{I}} : \mathbb{R}^n \rightarrow \mathbb{R}^{m_{\mathcal{I}}}$ are continuously differentiable

- ▶ Physics-constrained, resource-constrained, etc.
- ▶ Long history of algorithms (penalty, SQP, interior-point, etc.)
- ▶ Comprehensive theory (even with lack of constraint qualifications)
- ▶ Effective software (Ipopt, Knitro, LOQO, etc.)

Learning: Prediction function

Our aim is to determine a prediction function $p \in \mathcal{P}$, where \mathcal{P} is some family of functions, such that

$$p(a_j)$$

yields an accurate prediction corresponding to any given input feature vector a_j .

Learning: Prediction function, parameterized

For practicality, let us say that the family is parameterized by some vector x such that

$$p(a_j, x)$$

yields an accurate prediction corresponding to any given input feature vector a_j .

Learning: Supervised

In the context of supervised learning, we have known input-output pairs $\{(a_j, b_j)\}_{j=1}^{n_o}$, then

$$\min_{x \in \mathbb{R}^n} \frac{1}{n_o} \sum_{j=1}^{n_o} \ell(p(a_j, x), b_j)$$

becomes our empirical-loss training problem to determine the optimal parameter vector x .

Learning: Supervised and regularized

If, in addition, we aim to impose some structure on the solution x , then we may consider

$$\min_{x \in \mathbb{R}^n} \frac{1}{n_o} \sum_{j=1}^{n_o} \ell(p(a_j, x), b_j) + r(x)$$

where r is a *regularization* function.

Learning: Supervised and regularized

If, in addition, we aim to impose some structure on the solution x , then we may consider

$$\min_{x \in \mathbb{R}^n} \frac{1}{n_o} \sum_{j=1}^{n_o} \ell(p(a_j, x), b_j) + r(x)$$

where r is a *regularization* function. But is this the right approach for *informed* learning?

Learning: Supervised and informed with *soft* constraints

Added to the loss (e.g., mean-squared error or other data-fitting term), we might consider

$$\min_{x \in \mathbb{R}^n} \frac{1}{n_o} \sum_{j=1}^{n_o} \ell(p(a_j, x), b_j) + \frac{1}{n_c} \sum_{j=1}^{n_c} \phi(p(\tilde{a}_j, x), \dots, \tilde{b}_j)$$

where $\{(\tilde{a}_j, \tilde{b}_j)\}_{j=1}^{n_c}$ are some known input-output pairs and ϕ encodes known information.

Learning: Supervised and informed through layer design

Another viable approach is to embed information through the prediction function itself such that

$$\min_{x \in \mathbb{R}^n} \frac{1}{n_o} \sum_{j=1}^{n_o} \ell(\hat{p}(a_j, x), b_j)$$

ensures that information is enforced with every forward pass. (Expense?)

Learning: Supervised and informed with *hard* constraints

Back to the “original” family for p , how about imposing hard constraints during training, as in

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \quad & \frac{1}{n_o} \sum_{j=1}^{n_o} \ell(p(a_j, x), b_j) \\ \text{s.t.} \quad & \varphi(p(\tilde{a}_j, x), \dots, \tilde{b}_j) = 0 \text{ (or } \leq 0) \text{ for all } i \in \{1, \dots, n_c\} \end{aligned}$$

such that we restrict attention to functions that are informed implicitly?

Expected-loss training problems

For the sake of generality/generalizability, the expected-loss objective function is

$$\int_{\mathcal{A} \times \mathcal{B}} \ell(p(a, x), b) d\mathbb{P}(a, b) \equiv \mathbb{E}_\omega[F(x, \omega)] =: f(x).$$

One might consider various paradigms for imposing the constraints:

- ▶ expectation constraints
- ▶ (distributionally) robust constraints
- ▶ probabilistic (i.e., chance) constraints

For our recent work, we consider constraints whose values and derivatives can be computed:

$$c_{\mathcal{E}}(x) = 0 \quad \text{and} \quad c_{\mathcal{I}}(x) \leq 0$$

e.g., as in imposing a fixed set of constraints corresponding to a fixed set of sample data.

Physics-informed learning (e.g., PINNs)




Photo: Karniadakis et al.

Fair learning

Let

- ▶ A be a feature vector
- ▶ Z be a sensitive feature vector
- ▶ B be the output/label

Given a prediction function p and loss ℓ , the expected-loss minimization problem is

$$\min_{x \in \mathbb{R}^n} \mathbb{E} \left[p \left(\underbrace{\phi \left(\begin{bmatrix} A \\ Z \end{bmatrix}, x \right)}_{\hat{B}}, B \right) \right].$$

However, the resulting loss might not be fair between subgroups in the population.

- ▶ Various criteria related to fairness (e.g., demographic parity, equalized odds, equalized opportunity) leading to various measures (e.g., accuracy equality, disparate impact, etc.)
- ▶ For example, in binary classification, disparate impact may be expressed as the constraint

$$\mathbb{P}[\hat{B} = b | Z = 1] = \mathbb{P}[\hat{b} = b | Z = 0] \quad \text{for each } b \in \{-1, 1\}$$

Constrained optimization (stochastic algorithms)

Our approach (as a stepping stone to tackling more difficult settings) is to consider

$$\begin{array}{ll} \min_{x \in \mathbb{R}^n} & f(x), \quad \text{where } f(x) = \mathbb{E}_\omega[F(x, \omega)] \\ \text{s.t.} & c_{\mathcal{E}}(x) = 0 \\ & c_{\mathcal{I}}(x) \leq 0 \end{array}$$

- ▶ Classical applications under uncertainty, constrained DNN training, etc.
- ▶ Besides cases involving a deterministic equivalent...
- ▶ ... very few algorithms so far (mostly penalty methods)

Outline

Motivation

Stochastic SQP

Extensions

Conclusion

Equality-constrained setting (to start)

Consider the *equality-constrained* optimization problem:

$$\begin{array}{ll} \min_{x \in \mathbb{R}^n} & f(x), \quad \text{where } f(x) = \mathbb{E}_\omega[F(x, \omega)] \\ \text{s.t.} & c(x) = 0 \end{array}$$

What kind of algorithm do we want?

Need to establish what we want/expect from an algorithm.

Note: We are interested in the **fully stochastic** regime.[†]

We assume:

- ▶ Feasible methods are not tractable
- ▶ ... so no projection methods, Frank-Wolfe, etc.
- ▶ “Two-phase” methods are not effective
- ▶ ... so should not search for feasibility, then optimize.

Finally, want to use techniques that can generalize to diverse settings.

[†]Alternatively, see Na, Anitescu, Kolar (2021, 2022) and others

Stochastic gradient method (SG)

Stochastic approximation by Herbert Robbins and Sutton Monro (1951)



Sutton Monro, former Lehigh faculty member

Stochastic gradient (*not* descent)

Suppose $\nabla f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is Lipschitz continuous with constant L

Algorithm SG : Stochastic Gradient

-
- 1: choose an initial point $x_1 \in \mathbb{R}^n$ and step sizes $\{\alpha_k\} > 0$
 - 2: **for** $k \in \{1, 2, \dots\}$ **do**
 - 3: set $x_{k+1} \leftarrow x_k - \alpha_k g_k$, where $\mathbb{E}[G_k | \mathcal{F}_k] = \nabla f(X_k)$ and $\mathbb{E}[\|G_k - \nabla f(X_k)\|_2^2 | \mathcal{F}_k] \leq M$
 - 4: **end for**
-

Notation: $\{(x_k, g_k)\}$ is a realization of the stochastic process $\{(X_k, G_k)\}$ with filtration $\{\mathcal{F}_k\}$

Not a descent method! ... but *eventual descent in expectation*:

$$\begin{aligned} f(X_{k+1}) - f(X_k) &\leq \nabla f(X_k)^T (X_{k+1} - X_k) + \frac{1}{2} L \|X_{k+1} - X_k\|_2^2 \\ &= -\alpha_k \nabla f(X_k)^T G_k + \frac{1}{2} \alpha_k^2 L \|G_k\|_2^2 \\ \implies \mathbb{E}[f(X_{k+1}) | \mathcal{F}_k] - f(X_k) &\leq -\alpha_k \|\nabla f(X_k)\|_2^2 + \frac{1}{2} \alpha_k^2 L \mathbb{E}[\|G_k\|_2^2 | \mathcal{F}_k]. \end{aligned}$$

Markovian: In any run, x_{k+1} depends only on x_k and random choice at iteration k .

SG theory

Theorem SG

Since $\mathbb{E}[G_k | \mathcal{F}_k] = \nabla f(X_k)$ and $\mathbb{E}[\|G_k - \nabla f(X_k)\|_2^2 | \mathcal{F}_k] \leq M$ for all $k \in \mathbb{N}$:

$$\alpha_k = \frac{1}{L} \quad \Rightarrow \quad \mathbb{E} \left[\frac{1}{k} \sum_{j=1}^k \|\nabla f(X_j)\|_2^2 \right] = \mathcal{O}(M)$$

$$\alpha_k = \Theta \left(\frac{1}{k} \right) \quad \Rightarrow \quad \mathbb{E} \left[\frac{1}{\left(\sum_{j=1}^k \alpha_j \right)} \sum_{j=1}^k \alpha_j \|\nabla f(X_j)\|_2^2 \right] \rightarrow 0$$

$$\Rightarrow \liminf_{k \rightarrow \infty} \mathbb{E}[\|\nabla f(X_k)\|_2^2] = 0$$

SG illustration




Figure: SG with fixed step size (left) vs. diminishing step sizes (right)

Sequential quadratic optimization (SQP)

Consider

$$\begin{array}{l} \min_{x \in \mathbb{R}^n} f(x) \\ \text{s.t. } c(x) = 0 \end{array}$$

with $J \equiv \nabla c$ and H positive definite over $\text{Null}(J)$, two viewpoints:

$$\begin{bmatrix} \nabla f(x) + J(x)^T y \\ c(x) \end{bmatrix} = 0$$

or

$$\begin{array}{l} \min_{d \in \mathbb{R}^n} f(x) + \nabla f(x)^T d + \frac{1}{2} d^T H d \\ \text{s.t. } c(x) + J(x)d = 0 \end{array}$$

both leading to the same “Newton-SQP system”:

$$\begin{bmatrix} H_k & J_k^T \\ J_k & 0 \end{bmatrix} \begin{bmatrix} d_k \\ y_k \end{bmatrix} = - \begin{bmatrix} \nabla f(x_k) \\ c_k \end{bmatrix}$$

SQP illustration




Figure: Illustrations of SQP subproblem solutions

SQP with backtracking line search

Algorithm guided by merit function with **adaptive** parameter τ defined by

$$\phi(x, \tau) = \tau f(x) + \|c(x)\|_1$$

Algorithm SQP w/ line search

1: choose $x_1 \in \mathbb{R}^n$, $\tau_0 \in \mathbb{R}_{>0}$, $\eta \in (0, 1)$

2: **for** $k \in \{1, 2, \dots\}$ **do**

3: **compute step**: solve

$$\begin{bmatrix} H_k & J_k^T \\ J_k & 0 \end{bmatrix} \begin{bmatrix} d_k \\ y_k \end{bmatrix} = - \begin{bmatrix} \nabla f(x_k) \\ c_k \end{bmatrix}$$

4: **update merit parameter**: set τ_k to ensure

$$\phi'(x_k, \tau_k, d_k) \leq -\Delta q(x_k, \tau_k, \nabla f(x_k), d_k) \ll 0$$

5: **compute step size**: backtracking line search to ensure $x_{k+1} \leftarrow x_k + \alpha_k d_k$ yields

$$\phi(x_{k+1}, \tau_k) \leq \phi(x_k, \tau_k) - \eta \alpha_k \Delta q(x_k, \tau_k, \nabla f(x_k), d_k)$$

6: **end for**

Convergence theory

Assumption

- ▶ $f, c, \nabla f$, and J bounded and Lipschitz
- ▶ singular values of J bounded below (i.e., the LICQ)
- ▶ $u^T H_k u \geq \zeta \|u\|_2^2$ for all $u \in \text{Null}(J_k)$ for all $k \in \mathbb{N}$

Theorem

- ▶ $\{\alpha_k\} \geq \alpha_{\min}$ for some $\alpha_{\min} > 0$
- ▶ $\{\tau_k\} \geq \tau_{\min}$ for some $\tau_{\min} > 0$
- ▶ $\Delta q(x_k, \tau_k, \nabla f(x_k), d_k) \rightarrow 0$ implies optimality error vanishes, specifically,

$$\|d_k\|_2 \rightarrow 0, \quad \|c_k\|_2 \rightarrow 0, \quad \|\nabla f(x_k) + J_k^T y_k\|_2 \rightarrow 0$$

Toward stochastic SQP

- ▶ In a stochastic setting, line searches are (likely) intractable
- ▶ However, for ∇f and ∇c , may have Lipschitz constants L and Γ
- ▶ Step #1: Design an **adaptive** SQP method with

step sizes determined by Lipschitz constants

- ▶ Step #2: Design a **stochastic** SQP method based on this approach

SQP with adaptive step sizes

Algorithm SQP w/o line search

- 1: choose $x_1 \in \mathbb{R}^n$, $\tau_0 \in \mathbb{R}_{>0}$, $\eta \in (0, 1)$
 2: **for** $k \in \{1, 2, \dots\}$ **do**
 3: **compute step:** solve

$$\begin{bmatrix} H_k & J_k^T \\ J_k & 0 \end{bmatrix} \begin{bmatrix} d_k \\ y_k \end{bmatrix} = - \begin{bmatrix} \nabla f(x_k) \\ c_k \end{bmatrix}$$

- 4: **update merit parameter:** set τ_k to ensure

$$\phi'(x_k, \tau_k, d_k) \leq -\Delta q(x_k, \tau_k, \nabla f(x_k), d_k) \ll 0$$

- 5: **compute step size:** set

$$\hat{\alpha}_k \leftarrow \frac{2(1-\eta)\Delta q(x_k, \tau_k, \nabla f(x_k), d_k)}{(\tau_k L + \Gamma)\|d_k\|_2^2} \quad \text{and} \quad \tilde{\alpha}_k \leftarrow \hat{\alpha}_k - \frac{4\|c_k\|_1}{(\tau_k L + \Gamma)\|d_k\|_2^2}$$

- 6: **then**

$$\alpha_k \leftarrow \begin{cases} \hat{\alpha}_k & \text{if } \hat{\alpha}_k < 1 \\ 1 & \text{if } \tilde{\alpha}_k \leq 1 \leq \hat{\alpha}_k \\ \tilde{\alpha}_k & \text{if } \tilde{\alpha}_k > 1 \end{cases}$$

- 7: then set $x_{k+1} \leftarrow x_k + \alpha_k d_k$
 8: **end for**
-

Convergence theory: Nearly identical as for SQP w/ line search.

Stochastic SQP with adaptive step sizes

Algorithm : Stochastic SQP1: choose $x_1 \in \mathbb{R}^n$, $\tau_0 \in \mathbb{R}_{>0}$, $\{\beta_k\} \in (0, 1)$ 2: **for** $k \in \{1, 2, \dots\}$ **do**3: **compute step**: solve

$$\begin{bmatrix} H_k & J_k^T \\ J_k & 0 \end{bmatrix} \begin{bmatrix} d_k \\ y_k \end{bmatrix} = - \begin{bmatrix} g_k \\ c_k \end{bmatrix}$$

4: **update merit parameter**: set τ_k to ensure

$$\phi'(x_k, \tau_k, d_k) \leq -\Delta q(x_k, \tau_k, g_k, d_k) \ll 0$$

5: **compute step size**: set

$$\hat{\alpha}_k \leftarrow \frac{\beta_k \Delta q(x_k, \tau_k, g_k, d_k)}{(\tau_k L + \Gamma) \|d_k\|_2^2} \quad \text{and} \quad \tilde{\alpha}_k \leftarrow \hat{\alpha}_k - \frac{4\|c_k\|_1}{(\tau_k L + \Gamma) \|d_k\|_2^2}$$

6: **then**

$$\alpha_k \leftarrow \begin{cases} \hat{\alpha}_k & \text{if } \hat{\alpha}_k < 1 \\ 1 & \text{if } \tilde{\alpha}_k \leq 1 \leq \hat{\alpha}_k \\ \tilde{\alpha}_k & \text{if } \tilde{\alpha}_k > 1 \end{cases}$$

7: **then** $x_{k+1} \leftarrow x_k + \alpha_k d_k$ 8: **end for**Assume $\{g_k\}$ is a realization of $\{G_k\}$ with $\mathbb{E}[G_k | \mathcal{F}_k] = \nabla f(X_k)$ and $\mathbb{E}[\|G_k - \nabla f(X_k)\|_2^2 | \mathcal{F}_k] \leq M$

Fundamental lemma

Recall in the unconstrained setting that

$$\mathbb{E}[f(X_{k+1})|\mathcal{F}_k] - f(X_k) \leq -\alpha_k \|\nabla f(X_k)\|_2^2 + \frac{1}{2}\alpha_k^2 L \mathbb{E}[\|G_k\|_2^2|\mathcal{F}_k]$$

Lemma

For all $k \in \mathbb{N}$ one finds (before taking expectations)

$$\begin{aligned} & \phi(X_{k+1}, \mathcal{T}_{k+1}) - \phi(X_k, \mathcal{T}_k) \\ \leq & \underbrace{-\mathcal{A}_k \Delta q(X_k, \mathcal{T}_k, \nabla f(X_k), D_k^{\text{true}})}_{\mathcal{O}(\beta_k), \text{ "deterministic" }} + \underbrace{\frac{1}{2}\mathcal{A}_k \beta_k \Delta q(X_k, \mathcal{T}_k, G_k, D_k)}_{\mathcal{O}(\beta_k^2), \text{ stochastic/noise }} + \underbrace{\mathcal{A}_k \mathcal{T}_k \nabla f(X_k)^T (D_k - D_k^{\text{true}})}_{\text{ due to adaptive } \mathcal{A}_k} \end{aligned}$$

Good merit parameter behavior

Lemma

Let $\mathcal{E} :=$ event that $\{\mathcal{T}_k\}$ eventually remains constant at $\mathcal{T}' \geq \tau_{\min} > 0$. Then, for large k ,

$$\mathbb{E}_\omega[\mathcal{A}_k \mathcal{T}_k \nabla f(X_k)^T (D_k - D_k^{\text{true}}) | \mathcal{F}_k \cap \mathcal{E}] = \beta_k^2 \mathcal{T}' \mathcal{O}(\sqrt{M})$$

Theorem

Conditioned on \mathcal{E} , for large k , one finds

$$\beta_k = \Theta(1) \implies \mathbb{E} \left[\frac{1}{k} \sum_{j=1}^k \Delta q(X_j, \mathcal{T}', \nabla f(X_j), D_j^{\text{true}}) \right] = \mathcal{O}(M)$$

$$\beta_k = \Theta\left(\frac{1}{k}\right) \implies \mathbb{E} \left[\frac{1}{\left(\sum_{j=1}^k \beta_j\right)} \sum_{j=1}^k \beta_j \Delta q(X_j, \mathcal{T}', \nabla f(X_j), D_j^{\text{true}}) \right] \rightarrow 0$$

Good merit parameter behavior

Lemma

Let $\mathcal{E} :=$ event that $\{\mathcal{T}_k\}$ eventually remains constant at $\mathcal{T}' \geq \tau_{\min} > 0$. Then, for large k ,

$$\mathbb{E}_\omega[\mathcal{A}_k \mathcal{T}_k \nabla f(X_k)^T (D_k - D_k^{\text{true}}) | \mathcal{F}_k \cap \mathcal{E}] = \beta_k^2 \mathcal{T}' \mathcal{O}(\sqrt{M})$$

Theorem

Conditioned on \mathcal{E} , for large k , one finds

$$\beta_k = \Theta(1) \implies \mathbb{E} \left[\frac{1}{k} \sum_{j=1}^k (\|\nabla f(X_j) + \nabla c(X_j)^T Y_j^{\text{true}}\|_2 + \|c(X_j)\|_2) \right] = \mathcal{O}(M)$$

$$\beta_k = \Theta\left(\frac{1}{k}\right) \implies \mathbb{E} \left[\frac{1}{\left(\sum_{j=1}^k \beta_j\right)} \sum_{j=1}^k \beta_j (\|\nabla f(X_j) + \nabla c(X_j)^T Y_j^{\text{true}}\|_2 + \|c(X_j)\|_2) \right] \rightarrow 0$$

Poor merit parameter behavior

$\{\mathcal{T}_k\} \searrow 0$:

- ▶ cannot occur if $\|G_k - \nabla f(X_k)\|_2$ is bounded uniformly
- ▶ occurs with small probability if distribution of G_k has “small tails”

$\{\mathcal{T}_k\}$ remains too large:

- ▶ under a modest assumption, occurs with probability zero

Numerical results: (Matlab) <https://github.com/frankecurtis/StochasticSQP>

CUTE problems with noise added to gradients with different noise levels

- ▶ Stochastic SQP: 10^3 iterations
- ▶ Stochastic Subgradient: 10^4 iterations and tuned over 11 values of penalty parameter




Figure: Box plots for feasibility errors (left) and optimality errors (right).

Outline

Motivation

Stochastic SQP

Extensions

Conclusion

Summary

Since our original work, we have considered various extensions.

- ▶ stronger convergence guarantees (convergence in probability \rightarrow almost-sure convergence)
- ▶ convergence of Lagrange multiplier estimates
- ▶ relaxed constraint qualifications
- ▶ worst-case complexity guarantees
- ▶ generally constrained problems (with inequality constraints as well)
- ▶ interior-point methods
- ▶ iterative linear system solvers and inexactness

Almost-sure convergence of merit function value

Convergence of the algorithm is driven by the exact merit function

$$\phi_\tau(X) = \tau f(X) + \|c(X)\|$$

Reductions in a local model of ϕ_τ can be tied to a stationarity measure

$$\Delta q_\tau(X, \nabla f(X), H, D^{\text{true}}) \quad \sim \quad \|\nabla f(X) + \nabla c(X)Y\|^2 + \|c(X)\|$$

Lemma

Suppose $\mathbb{E}[G_k | \mathcal{F}_k] = \nabla f(X_k)$ and $\mathbb{E}[\|G_k - \nabla f(X_k)\|^2 | \mathcal{F}_k] \leq M$. Then, by a classical theorem of Robbins and Siegmund (1971), one finds that, almost surely,

$$\lim_{k \rightarrow \infty} \{\phi_\tau(X_k)\} \text{ exists and is finite and}$$
$$\liminf_{k \rightarrow \infty} \Delta q_\tau(X_k, \nabla f(X_k), H_k, D_k^{\text{true}}) = 0$$

Almost-sure convergence of the primal iterates

If $\{X_k\}$ stays within a neighborhood of x_* almost surely, where x_* is a stationary point at which a generalization of the Polyak–Lojasiewicz condition holds, then almost-sure convergence follows:

Theorem

Suppose that there exists $x_* \in \mathcal{X}$ with $c(x_*) = 0$, $\mu \in \mathbb{R}_{>1}$, and $\epsilon \in \mathbb{R}_{>0}$ such that for all

$$x \in \mathcal{X}_{\epsilon, x_*} := \{x \in \mathcal{X} : \|x - x_*\|_2 \leq \epsilon\}$$

one finds that

$$\phi_\tau(x) - \phi_\tau(x_*) \begin{cases} = 0 & \text{if } x = x_* \\ \in (0, \mu(\tau\|Z(x)^T \nabla f(x)\|_2^2 + \|c(x)\|_2)] & \text{otherwise,} \end{cases}$$

where for all $x \in \mathcal{X}_{\epsilon, x_*}$ one defines $Z(x) \in \mathbb{R}^{n \times (n-m)}$ as some orthonormal matrix whose columns form a basis for the null space of $\nabla c(x)^T$. Then, if $\limsup_{k \rightarrow \infty} \{\|X_k - x_*\|_2\} \leq \epsilon$ almost surely, it follows that

$$\{\phi_\tau(X_k)\} \xrightarrow{a.s.} \phi_\tau(x_*), \quad \{X_k\} \xrightarrow{a.s.} x_*, \quad \text{and} \quad \left\{ \begin{bmatrix} \nabla f(X_k) + \nabla c(X_k) Y_k^{\text{true}} \\ c(X_k) \end{bmatrix} \right\} \xrightarrow{a.s.} 0.$$

Lagrange multiplier convergence

Theorem

Suppose (x_*, y_*) is a stationary point. Then, for any $k \in \mathbb{N}$, one finds $\|X_k - x_*\|_2 \leq \epsilon$ implies

$$\|Y_k - y_*\|_2 \leq \kappa_y \|X_k - x_*\|_2 + r^{-1} \|\nabla f(X_k) - G_k\|_2$$


and $\|Y_k^{\text{true}} - y_*\|_2 \leq \kappa_y \|X_k - x_*\|_2$ for some $(\kappa, r) \in \mathbb{R}_{>0} \times \mathbb{R}_{>0}$.

Computed multipliers *always* have error. Consider *averaged* multipliers $\{Y_k^{\text{avg}}\}$ instead.

Theorem

If the iterate sequence converges almost surely to x_* , i.e., $\{X_k\} \xrightarrow{\text{a.s.}} x_*$, then

$$\{Y_k^{\text{true}}\} \xrightarrow{\text{a.s.}} y_* \quad \text{and} \quad \{Y_k^{\text{avg}}\} \xrightarrow{\text{a.s.}} y_*.$$

Constrained logistic regression: **australian** dataset (LIBSVM)

Relaxing constraint qualifications

Use a step decomposition method, handled infeasible and/or degenerate problems as well.




Figure: Box plots for feasibility errors (left) and optimality errors (right).

Complexity of $\mathcal{O}(\epsilon^{-2})$ for deterministic algorithm

All reductions in the merit function can be cast in terms of smallest τ .

Since τ_{\min} is determined by the initial point, *it will be reached.*

Theorem

For any $\epsilon \in (0, 1)$, there exists $(\kappa_1, \kappa_2) \in (0, \infty) \times (0, \infty)$ such that

$$\|\nabla f(x_k) + J_k^T y_k\| \leq \epsilon \text{ and } \sqrt{\|c_k\|_1} \leq \epsilon$$

in a number of iterations no more than

$$\left(\frac{\tau_0(f_1 - f_{\inf}) + \|c_1\|_1}{\min\{\kappa_1, \kappa_2 \tau_{\min}\}} \right) \epsilon^{-2}.$$

Complexity of $\tilde{\mathcal{O}}(\epsilon^{-4})$ for stochastic algorithm

Theorem

Suppose the algorithm is run k_{\max} iterations with

- ▶ $\beta_k = \gamma/\sqrt{k_{\max} + 1}$ and
- ▶ the merit parameter is reduced at most $s_{\max} \in \{0, 1, \dots, k_{\max}\}$ times.

Let K_* be sampled uniformly over $\{1, \dots, k_{\max}\}$. Then, with probability $1 - \delta$,

$$\mathbb{E}[\|\nabla f(X_{K_*}) + J(X_{K_*})^T Y_{k_*}^{\text{true}}\|_2^2 + \|c(X_{K_*})\|_1] \leq \frac{\tau_0(f(x_1) - f_{\text{inf}}) + \|c(x_1)\|_1 + M}{\sqrt{k_{\max} + 1}} + \frac{(\tau_{-1} - \tau_{\min})(s_{\max} \log(k_{\max}) + \log(1/\delta))}{\sqrt{k_{\max} + 1}}$$

Theorem



If the stochastic gradient estimates are sub-Gaussian, then w.p. $1 - \bar{\delta}$

$$s_{\max} = \mathcal{O}\left(\log\left(\log\left(\frac{k_{\max}}{\bar{\delta}}\right)\right)\right).$$

Inequality-constrained optimization

Stochastic SQP for inequality constrained problems

- ▶ employed in an ϵ -constraint method for fair machine learning



$$\min_{x \in \mathbb{R}^n} \frac{1}{N_o} \sum_{(v_i, y_i) \in D_o} \ell(x, v_i, y_i) \quad \text{s.t.} \quad -\epsilon \leq \frac{1}{N_c} \sum_{(v_i, a_i) \in D_c} (a_i - \bar{a}) x^T v_i \leq \epsilon$$

Interior-point methods

Stochastic *single-loop* algorithm (prescribed barrier sequence $\{\mu_k\} \searrow 0$) with convergence guarantees.




Figure: Deterministic setting (left) and stochastic setting (right)

Iterative methods and inexactness

Inexact subproblem solves

- ▶ stochasticity and inexactness(!)

Iterative methods employed to solve

$$\begin{bmatrix} H_k & J_k^T \\ J_k & 0 \end{bmatrix} \begin{bmatrix} d_k \\ y_k \end{bmatrix} = - \begin{bmatrix} g_k \\ c_k \end{bmatrix}$$

termination tests to determine when an inexact solution is sufficient for convergence.

Physics-informed learning





Figure: True solution (left) and predicted solutions (right).

Outline

Motivation

Stochastic SQP

Extensions

Conclusion

Summary

Consider stochastic-gradient-based algorithms for solving problems of the form:

$$\begin{array}{l} \min_{x \in \mathbb{R}^n} f(x), \quad \text{where } f(x) = \mathbb{E}_\omega[F(x, \omega)] \\ \text{s.t. } c_{\mathcal{E}}(x) = 0 \\ \quad \quad c_{\mathcal{I}}(x) \leq 0 \end{array}$$


Equality-constraints-only setting:

- ▶ convergence in probability with complexity guarantees
- ▶ almost-sure convergence of primal iterates and averaged Lagrange multipliers
- ▶ relaxed constraint qualifications
- ▶ inexact subproblem solves

Generally constrained setting (with inequality constraints as well):

- ▶ stochastic SQP
- ▶ stochastic interior-point (bounds only so far, but generally constrained in progress)

Collaborators and references



- ▶ A. S. Berahas, F. E. Curtis, D. P. Robinson, and B. Zhou, “Sequential Quadratic Optimization for Nonlinear Equality Constrained Stochastic Optimization,” *SIAM Journal on Optimization*, 31(2):1352–1379, 2021.
- ▶ A. S. Berahas, F. E. Curtis, M. J. O’Neill, and D. P. Robinson, “A Stochastic Sequential Quadratic Optimization Algorithm for Nonlinear Equality Constrained Optimization with Rank-Deficient Jacobians,” to appear in *Mathematics of Operations Research*.
- ▶ F. E. Curtis, D. P. Robinson, and B. Zhou, “Inexact Sequential Quadratic Optimization for Minimizing a Stochastic Objective Subject to Deterministic Nonlinear Equality Constraints,” <https://arxiv.org/abs/2107.03512>.
- ▶ F. E. Curtis, M. J. O’Neill, and D. P. Robinson, “Worst-Case Complexity of an SQP Method for Nonlinear Equality Constrained Stochastic Optimization,” to appear in *Mathematical Programming*.
- ▶ F. E. Curtis, S. Liu, and D. P. Robinson, “Fair Machine Learning through Constrained Stochastic Optimization and an ϵ -Constraint Method,” to appear in *Optimization Letters*.
- ▶ F. E. Curtis, D. P. Robinson, and B. Zhou, “Sequential Quadratic Optimization for Stochastic Optimization with Deterministic Nonlinear Inequality and Equality Constraints,” <https://arxiv.org/abs/2302.14790>.
- ▶ F. E. Curtis, V. Kungurtsev, D. P. Robinson, and Q. Wang, “A Stochastic-Gradient-based Interior-Point Algorithm for Solving Smooth Bound-Constrained Optimization Problems,” <https://arxiv.org/abs/2304.14907>.